

## PRINTED REFERENCES

- Batcher, M.S. and S.A. Stiles. 2000. Element stewardship abstract for *Lonicera maackii* (Amur honeysuckle), *Lonicera morrowii* (Morrow's honeysuckle), *Lonicera tatarica* (Tatarian honeysuckle), and *Lonicera x bella* (Bell's honeysuckle): The bush honeysuckles. The Nature Conservancy, Arlington, VA.
- Brock, T.D. 2004. Use of herbicide in the eradication of honeysuckle (Wisconsin). *Ecological Restoration* 22(2):145-146.
- Cochrane, T.S. 1995. *Lonicera maackii* (Caprifoliaceae) naturalized in Wisconsin. *The Michigan Botanist* 34:79-83.
- Czarapata, E.J. 2005. Exotic Honeysuckles. p. 32-35. In: *Invasive Plants of the Upper Midwest: An Illustrated Guide to Their Identification and Control*. University of Wisconsin Press, Madison, WI.
- Glass, S. 1994. Experiment finds less herbicide needed to control buckthorn. *Restoration and Management Notes*. 12(1):93.
- Gould, A.M.A. and D.L. Gorchov. 2000. Effects of the exotic invasive shrub *Lonicera maackii* on the survival and fecundity of three species of native annuals. *American Midland Naturalist*. 144:36-50.
- Hoffman, R. and K. Kearns (eds.). 1998. Exotic bush honeysuckles. p. 19-21. In: *Wisconsin manual of control recommendations for ecologically invasive plants*. Bureau of Endangered Resources, Wisconsin Department of Natural Resources, Madison, WI.
- Luken, J.O. 1996. *Lonicera maackii*, *L. morrowii*, *L. tatarica*: Bush honeysuckle. p. 60-61. In: J.M. Randall and J. Marinelli (eds.). *Invasive plants: Weeds of the global garden*. Brooklyn Botanic Garden Publications, Handbook #149, Brooklyn, NY.
- Schmidt, K.A. and C.J. Whelan. 1999. Effects of exotic *Lonicera* and *Rhamnus* on songbird nest predation. *Conservation Biology* 13(6):1502-1506.

## WEBSITES

- [www.nps.gov/plants/alien/factmain.htm](http://www.nps.gov/plants/alien/factmain.htm)  
Plant Conservation Alliance Alien Plant Working Group fact sheets on alien plant invaders of natural areas, including honeysuckle
- [www.ipaw.org/invasives/honeysuckle/index.htm](http://www.ipaw.org/invasives/honeysuckle/index.htm)  
Invasive Plants Association of Wisconsin summary on invasive honeysuckles
- <http://plants.usda.gov>  
U.S.D.A. Natural Resources Conservation Service summary of United States invasive weeds
- [www.dnr.state.mn.us/invasives/terrestrialplants/woody/exotichoneysuckles.html](http://www.dnr.state.mn.us/invasives/terrestrialplants/woody/exotichoneysuckles.html)  
Minnesota Department of Natural Resources summary of honeysuckles
- [www.nrc.ca/scf.ec.gc.ca/publications/inv/14\\_e.cfm\\*targ6](http://www.nrc.ca/scf.ec.gc.ca/publications/inv/14_e.cfm*targ6)  
Environment Canada, Canadian Wildlife Service publication of invasive plants of natural habitats in Canada
- [www.botany.wisc.edu/wisflora](http://www.botany.wisc.edu/wisflora)  
University of Wisconsin-Madison Herbarium checklist of the vascular plants of Wisconsin; shows range of honeysuckles in Wisconsin
- [www.dnr.state.wi.us/invasives/plants.asp](http://www.dnr.state.wi.us/invasives/plants.asp)  
Wisconsin Dept. of Natural Resources Endangered Resources; fact sheets on invasives including honeysuckles
- [www.imapinvasives.org/GIST/ESA/esapages/loni\\_spp.html](http://www.imapinvasives.org/GIST/ESA/esapages/loni_spp.html)  
The Nature Conservancy summary of honeysuckles

## UW-EXTENSION COUNTY OFFICES

- |                           |                           |
|---------------------------|---------------------------|
| Adams: 608-339-4237       | Marathon: 715-261-1230    |
| Ashland: 715-682-7017     | Marinette: 715-732-7510   |
| Barron: 715-537-6250      | Marquette: 608-297-3141   |
| Bayfield: 715-373-6104    | Menominee: 715-799-4654   |
| Brown: 920-391-4610       | Milwaukee: 414-256-4644   |
| Buffalo: 608-685-6256     | Monroe: 608-269-8722      |
| Burnett: 715-349-2151     | Oconto: 920-834-6845      |
| Calumet: 920-849-1450     | Oneida: 715-365-2750      |
| Chippewa: 715-726-7950    | Outagamie: 920-832-5121   |
| Clark: 715-743-5121       | Ozaukee: 262-238-8288     |
| Columbia: 608-742-9680    | Pepin: 715-672-5214       |
| Crawford: 608-326-0223    | Pierce: 715-273-6781      |
| Dane: 608-224-3700        | Polk: 715-485-8600        |
| Dodge: 920-386-3790       | Portage: 715-346-1316     |
| Door: 920-746-2260        | Price: 715-339-2555       |
| Douglas: 715-395-1363     | Racine: 262-886-8460      |
| Dunn: 715-232-1636        | Richland: 608-647-6148    |
| Eau Claire: 715-839-4712  | Rock: 608-757-5696        |
| Florence: 715-528-4480    | Rusk: 715-532-2151        |
| Fond du Lac: 920-929-3170 | St. Croix: 715-684-3301   |
| Forest: 715-478-7797      | Sauk: 608-355-3250        |
| Grant: 608-723-2125       | Sawyer: 715-634-4839      |
| Green: 608-328-9440       | Shawano: 715-526-6136     |
| Green Lake: 920-294-4032  | Sheboygan: 920-459-5900   |
| Iowa: 608-935-0391        | Taylor: 715-748-3327      |
| Iron: 715-561-2695        | Trempealeau: 715-538-2311 |
| Jackson: 715-284-4257     | Vernon: 608-637-5276      |
| Jefferson: 920-674-7295   | Vilas: 715-479-3648       |
| Juneau: 608-847-9329      | Walworth: 262-741-4951    |
| Kenosha: 262-857-1945     | Washburn: 715-635-4444    |
| Kewaunee: 920-388-7141    | Washington: 262-335-4477  |
| La Crosse: 608-785-9593   | Waukesha: 262-548-7770    |
| Lafayette: 608-776-4820   | Waupaca: 715-258-6230     |
| Langlade: 715-627-6236    | Waushara: 920-787-0416    |
| Lincoln: 715-539-1072     | Winnebago: 920-232-1970   |
| Manitowoc: 920-683-4168   | Wood: 715-421-8440        |

©2009 by the Board of Regents of the University of Wisconsin System doing business as the division of Cooperative Extension of the University of Wisconsin Extension.

An EEO/Affirmative Action employer, University of Wisconsin Extension provides equal opportunities in employment and programming, including Title IX and ADA requirements.

This brochure was produced by Dr. Laura Jull, UW-Madison Woody Ornamental Extension Specialist, with layout by Annika Sargent, UW-Extension Environmental Resources Center. This brochure was produced by the Urban Horticulture-Wisconsin Program of the USDA CSREES, grant number 2002-45060-01394.

Additional honeysuckle brochures are available from your county Extension office and local Wisconsin DNR Service Center (PUB-FR-448-2009-Invasive Exotic Shrub Honeysuckles).

Reviewed by: Dr. Joe Neal, Weed Specialist, Dept. of Horticultural Science, N.C. State University, Raleigh, NC; Dr. Brian Hudelson, Plant Disease Diagnostician, Dept. of Plant Pathology, UW-Madison, Madison, WI; Kelly Kearns, Bureau of Endangered Resources, Wisconsin Department of Natural Resources, Madison, WI; Steve Glass, University of Wisconsin-Madison Arboretum Naturalist, Madison, WI; Paul Hartman, former Horticulture County Agent, Brown County UW-Extension, Green Bay, WI.



# Invasive Exotic Shrub HONEYSUCKLES



## MAJOR THREATS TO MIDWESTERN WOODLANDS

## THREATS

Invasive exotic shrub honeysuckles

- ❖ can displace native understory vegetation;
- ❖ form an impenetrable understory layer;
- ❖ may degrade wildlife habitat;
- ❖ can cause long-term decline of forests by shading out other woody and herbaceous plants;
- ❖ may compete with native plants for pollinators, reducing seed set of native plants;
- ❖ may not serve as a source of high-fat, nutrient-rich fruit for migrating birds.

## IDENTIFICATION

**Growth:** Invasive exotic shrub honeysuckles are medium to large-sized, deciduous, multi-stemmed shrubs growing 6–15 feet tall and 6–12 feet wide. They have dense, upright, vase-shaped to rounded forms. Amur honeysuckle (*Lonicera maackii*) and Bell's honeysuckle (*L. x bella*), usually grow slightly taller than Morrow's honeysuckle (*L. morrowii*) and Tatarian honeysuckle (*L. tatarica*). Honeysuckles are fast growing and shallow rooted, forming dense colonies when they invade woodlands.

**Leaves:** Leaves of invasive exotic shrub honeysuckles are simple, opposite, oval to oblong, with entire margins and short petioles. They typically measure 1–2½ inches long and 1–1½ inches wide. Amur honeysuckle has darker, larger, somewhat glossy leaves (2–3 inches long) that taper to a distinct, long, slender point at the tip and are slightly hairy. Morrow's honeysuckle has hairy, gray-green leaves, while Tatarian honeysuckle has smooth, hairless, dull, bluish-green leaves. Bell's honeysuckle, a hybrid between Tatarian and Morrow's honeysuckle, has

dull leaves that are slightly hairy on the undersides. Invasive exotic shrub honeysuckles are easy to find in early spring and late autumn, as they tend to leaf out early and retain their leaves late into the fall. Invasive exotic shrub honeysuckles do not have distinctive fall color.

**Stems:** Stems of older, invasive exotic shrub honeysuckles are often hollow, with light, grayish-brown, shaggy bark in long strips. Buds along the stem are opposite and project at almost right angles to the stem. In comparison, native shrub honeysuckle stems have solid, white piths (i.e., stem centers).

**Flowers:** Invasive exotic shrub honeysuckles flower from early to late May until early June, depending on location. Flowers are borne in pairs at the nodes along the stem (i.e., in the leaf axils). Flowers are generally less than 1 inch long, fragrant, tubular, two-lipped and five-lobed, with five stamens (i.e., male floral parts). Nectar collects at the base of the flower tube and is attractive to both insect and hummingbird pollinators.

Amur and Morrow's honeysuckles produce white flowers that fade to yellow with age but Morrow's honeysuckle flowers are hairy and are borne on hairy peduncles (i.e., flower stems). Tatarian honeysuckle produces smooth flowers that are white, pink or crimson-red (especially on cultivars)



and do not yellow with age. Bell's honeysuckle has white to pink flowers, fading to yellow, on slightly hairy stems. *Lonicera* 'Freedom' and *L.* 'Honeyrose', two cultivars that are the result of a complex hybrid of several honeysuckle species, produce pink-tinged white flowers and rosy-red flowers, respectively.

**Fruit and Seed:** Honeysuckle fruit ranges from red to orange-yellow, depending on species and cultivar, and ripen in mid-summer (Morrow's, Tatarian, Bell's, Freedom, and Honeyrose honeysuckles) or early to mid autumn (Amur honeysuckle). The fleshy, ¼-inch diameter berries contain many seeds. Seeds are less than ⅛-inch long, oval, flattened, and yellow. Every year, invasive exotic shrub honeysuckles produce large quantities of viable seed that are readily dispersed by birds. The seeds require a cold, moist period to break dormancy, which occurs in the soil during winter. The germination rate of honeysuckle seeds is high under a wide range of environmental conditions. However, germination is highest on open ground or where understory development is sparse. Honeysuckle seeds remain viable in soil for two or more years.

## HISTORY


Invasive exotic honeysuckles are native to Asia and southern Russia. They were introduced into North America as ornamentals in the mid-18<sup>th</sup> and 19<sup>th</sup> centuries, due to their showy flowers and fruit. They were also used for wildlife food and cover, and soil erosion control. Bell's honeysuckle (*L. x bella*), a hybrid of Tatarian honeysuckle (*L. tatarica*) and Morrow's honeysuckle (*L. morrowii*), has quickly become as invasive as its parents. Unfortunately, some invasive exotic honeysuckles, especially the Tatarian honeysuckle cultivars 'Arnold Red' and 'Zabelii', as well as Freedom and Honeyrose honeysuckles, are still sold due to their ornamental characteristics and form, but should be considered invasive and should not be planted.

## DISTRIBUTION AND HABITAT

Invasive exotic honeysuckles range from southern New England and Canada south into North Carolina, west to the Great Plains and north into the Midwest. They often invade woodlands, especially those that are grazed or disturbed. They can also occur along lakeshores, forest edges, abandoned fields, pastures, roadsides, and other open, upland habitats. Morrow's and Bell's honeysuckle can also invade sand plains, bogs and fens. These honeysuckles can live under a broad range of light and moisture conditions, as well as in many different plant communities. However, they do not perform as well in shady environments. Large, urban areas are often invaded by honeysuckles. However, rural infestations have occurred where honeysuckles have been introduced to provide wildlife cover and food.



Invasive exotic honeysuckles have been used as a hedge shrub due to their ornamental characteristics and form, but should not be used due to their invasiveness.



Invasive exotic honeysuckles invade forest understories, crowding out native shrubs, groundcovers, and ephemerals.

## SIMILAR SPECIES

Other exotic honeysuckles may look like their invasive cousins, but they are not considered invasive plants. These include Clavey's Dwarf and Mini Globe honeysuckles (*L. x xylosteoides* 'Clavey's Dwarf' and 'Mini Globe'), and Emerald Mound honeysuckle (*L. xylosteum* 'Emerald Mound'). Other native, rarely seen shrub honeysuckles include American fly honeysuckle (*L. canadensis*), bearberry honeysuckle (*L. involucrata*), swamp fly honeysuckle (*L. oblongifolia*), and mountain fly honeysuckle (*L. villosa*). Except for the native swamp fly honeysuckle, exotic shrub honeysuckles have hairy styles (i.e., female reproductive structures). Bush honeysuckle (*Diervilla lonicera*), is not a true honeysuckle, but is related, native, and has small, yellow flowers at the tips of branches in summer, and dry, brown capsules in late summer.

Some honeysuckles are twining, woody vines, rather than shrubs. Native examples include the grape honeysuckle (*L. reticulata*, formerly *L. prolifera*); hairy honeysuckle, (*L. hirsuta*), and limber honeysuckle (*L. dioica*). Non-native, vine honeysuckles, such as Dropmore Scarlet honeysuckle (*L. x brownii* 'Dropmore Scarlet'), yellow honeysuckle (*L. flava*), goldflame honeysuckle (*L. x heckrottii*),

Mandarin honeysuckle (*L. 'Mandarin'*), woodbine honeysuckle (*L. periclymenum*), and trumpet honeysuckle (*L. sempervirens*), are not considered invasive. However, the non-native Japanese honeysuckle vine (*L. japonica*), is highly invasive and readily grows in native areas, especially woodlands. This honeysuckle has fragrant, white flowers that turn yellow with age, and produces black fruit. This species is very invasive in the southern Midwest and throughout the eastern U.S.

## SPREAD

Invasive exotic honeysuckles can rapidly form dense shrub layers in the forest understory. These honeysuckles produce large numbers of fruit that are highly attractive to birds. In the eastern United States, over twenty species of birds feed on honeysuckle fruit. Thus, birds commonly move honeysuckle seeds across the landscape. Invasive exotic honeysuckle seedlings grow in areas with sparse vegetation, especially under tall trees and shrubs. They also spread vegetatively by producing suckers and sprouts at the base of the plant, especially after severe pruning. Because of this, invasive exotic honeysuckles tend to persist in an area once they have become established and can not be easily removed by cutting alone.

Invasive exotic honeysuckles are responsible for crowding and shading out many native trees, shrubs, groundcovers, and spring wildflowers. In addition, they may compete for pollinators, reducing fruit formation and seed set of native species. Invasive exotic honeysuckles may also have a negative impact on wildlife. American robins (*Turdus migratorius*) nesting in invasive shrub honeysuckles experienced higher predation than those nesting in native species, due to lower nest heights, a lack of protective thorns, and a more conducive branch structure for predator movement. However, these results were specific to a single location and it is not known whether they will be applicable to other sites or bird species. Finally, honeysuckle fruit, while plentiful, may not offer migrating birds the high-fat, nutrient-rich food source they need for long flights.

## CONTROL METHODS

Control of invasive exotic honeysuckles is best achieved with early identification, and removal of isolated plants before they begin to produce seed. Once established, honeysuckles can shade out existing vegetation and prevent establishment of the native understory. In large infestations of honeysuckle, larger, seed-producing plants should be removed first.

**Hand pulling:** Honeysuckle plants with a stem diameter of ½ inch or less can be easily removed by hand pulling when soil is moist. Because honeysuckles have shallow roots, larger plants can be dug or pulled out using a rope or chain placed around the base of the plant. All larger roots must be removed as the shrubs can resprout from any remaining roots. Digging will disturb the soil, which can lead to honeysuckle reinvasions. Be sure to revisit the site the following summer to remove newly emerged plants.

**Fire:** Prescribed burns in early spring or autumn show promise in controlling invasive honeysuckle seedlings, but kills only the tops of larger shrubs, which can resprout. Fire works best on seedlings in fire-adapted plant communities (e.g., prairies, savannas, and oak woods), and should not be used where native plant communities might be adversely affected. Repeated burning every one to two years may be necessary. Consult with a restoration expert before attempting a burn. When burning, be sure to follow all local ordinances and state fire codes, and obtain local permits as needed.

**Cutting:** Larger plants that are hard to remove by hand pulling should be cut at the base with a lopper, hand saw, or carefully with a chainsaw. Shrubs can resprout from cut stumps or from the root systems if not treated with herbicides. Repeated basal pruning during the growing season may eventually weaken the plant resulting in reduced sprouting.

**Herbicides:** The effectiveness of cutting can be improved by immediately painting or spraying

cut stumps with a selective herbicide such as triclopyr (e.g., Ortho® Brush-B-Gon® or Garlon 4®) or a non-selective herbicide such as glyphosate. Glyphosate is the active ingredient in many herbicides including Roundup Pro®, Touchdown®, or Rodeo® (for use near waterways). Use a 20–25% (by volume) solution to ensure death of the plant and to prevent resprouting. Apply the herbicide to stumps immediately after cutting using a low-pressure hand sprayer, spray bottle, or sponge applicator. Herbicide applications to stumps should be directed to the vascular (i.e., water-conducting) tissue located just inside of the bark. Follow-up treatments may be needed for the next several years if resprouting occurs.

For younger, smaller shrubs, a basal bark application of a 12.5% (by volume) oil-formulated triclopyr solution (Garlon 4®) may be used to control honeysuckle in autumn without the need for cutting. Oil formulations of triclopyr are more effective than triclopyr formulations diluted in water. Triclopyr is a selective herbicide and only kills broad-leaved plants. It does not harm most grasses. This herbicide is volatile and should not be used at temperatures above 80°F.

Foliar applications of herbicides to seedlings and larger plants can also be effective. However, this technique is best reserved for treating areas with large numbers of honeysuckle seedlings, as non-target vegetation can be easily damaged or killed. Herbicide concentrations of triclopyr or glyphosate are generally lower for foliar sprays (1–3% by volume) than for stump applications.

Dyes can be added to the herbicide formulation to help identify treated areas. As with any pesticide, read the label prior to use to verify that the rates listed here are consistent with those legally allowed for on the label, and to read about proper safety precautions.

Because invasive honeysuckles retain their leaves and continue to grow into late autumn, the best

time for foliar sprays of glyphosate to treat honeysuckles is in mid to late autumn when many non-target plants are going dormant and are least likely to be damaged. In addition, in autumn, honeysuckle shrubs are transporting nutrients to their roots and thus uptake of herbicides is best at this time resulting in the highest mortality. Winter applications are also very successful for fresh cut stumps, decreasing the risk of damaging non-target species. However, if shrubs are cut in winter and herbicide treatments are not applied, vigorous resprouting occurs in spring. Spring cut-stump applications of triclopyr after budbreak, can effectively control honeysuckles.

Repetition of mechanical and chemical control methods may be necessary for at least three to five years in order to deplete honeysuckle plants and their seed bank. Replant areas that were infested with invasive species with native species tolerant to existing environmental conditions. This can help prevent reinvasion of invasive species.

**Biological Control:** At this time, no biological control agents are available to control any of the exotic shrub honeysuckles.

**Education:** One of the best honeysuckle control methods is education. Tell your neighbors about invasive honeysuckles. A neighbors' honeysuckle can produce large amounts of fruit and seed that can be disseminated into your backyard and the surrounding neighborhood by birds. Encourage your neighbors to remove their invasive honeysuckles and monitor their yards for seedlings.

**NOTE:** References to pesticide and other products in this publication are for your convenience and are not an endorsement or criticism of one product over similar products. You are responsible for using pesticides according to the manufacturer's current label directions. Follow directions exactly to protect the environment and people from pesticide exposure. Failure to do so violates the law.